

Acoustic peaks in the CMB: a matter of standard causal boundary conditions on primordial density anisotropies

David H. Oaknin

Department of Physics, Technion, Haifa, 32000, Israel.

e-mail: doaknin@physics.technion.ac.il

The pattern of acoustic peaks in the sub-horizon power spectrum of primordial density anisotropies at recombination can be naturally understood in the framework of standard Friedmann-Robertson-Walker cosmology (without inflation) as a consequence of the boundary conditions imposed by the causal horizon on the statistical two-points correlation functions: the sub-horizon spectrum is discrete (harmonic), with comoving modes located at $k_n = n \frac{\pi}{H_{eq}^{-1}}$, $n = 1, 2, \dots$, because the causally connected patch of the universe at recombination is compact, with comoving radius H_{eq}^{-1} . The results presented in this paper complement those presented in [1], where it was shown that the scale invariance of the primordial density anisotropies over comoving scales of cosmological size is also a consequence of the boundary conditions imposed by causality. Together these results lay an appealing theoretical alternative to the inflationary paradigm as the ultimate answer for the origin of cosmological structures in standard cosmology.

The extremely uniform blackbody radiation at $T \sim 2.7$ K that permeates the whole visible universe [2] is a relic signature of the hotter and denser earlier stage at $T_{eq} \sim 3 \times 10^3$ K when free electrons and protons recombined into atomic hydrogen and the electromagnetic radiation decoupled from the thermal cosmic plasma [3]. The anisotropies observed over the sky $\frac{\Delta T(\theta, \phi)}{T} \lesssim 10^{-5}$ in the local temperature of this blackbody radiation [4], known commonly as the cosmic microwave background (CMB), are the imprint left by tiny random primordial density fluctuations in the, otherwise, homogeneous and isotropic cosmic plasma at the instant of recombination.

Precision measurements of the tiny anisotropies in the temperature of the CMB over the sky have become an extremely powerful tool in today cosmology. Yet the theoretical

understanding of the ultimate mechanism that seeded the primordial density anisotropies at the instant of recombination is unclear. It is widely believed that the statistical correlations of these primordial anisotropies cannot be explained by causal physics in the framework of the, otherwise, extraordinarily successful standard Friedmann-Robertson-Walker (FRW) cosmology. The main reason for this belief is, in vague words, that primordial anisotropies extend over comoving volumes of cosmological size, which are much larger than the comoving causal horizon at the instant of decoupling H_{eq}^{-1} . The puzzling apparent contradiction is known in the literature as the *origin of structures problem* of standard cosmology. It is closely related to the so-called *horizon problem* and has become the source of many speculative ideas in cosmology for, at least, the last twenty years.

The most outstanding of the proposals that have been considered to solve the *origin of structures problem* and *horizon problem* of standard cosmology is the so-called inflationary paradigm. The paradigm assumes that the patch of the universe visible at present was once causally connected in the remote past, at $T \lesssim T_{Planck} \sim 10^{28}$ eV, and primordial density anisotropies were imprinted then. A finite period of exponential expansion, the inflation, would have supposedly stretched that diminute comoving patch, beyond the causal horizon, to the size of the currently visible universe. The inflation is regarded today as the natural solution to the problems of standard cosmology, even though all the attempts to implement it in models of particle cosmology have failed.

In reference [1] we showed that the analysis that led to formulate the *origin of structures problem* and *horizon problem* of standard cosmology were based on an unjustified, and erroneous, approximation that had gone unnoticed. We indeed showed that the generic statistical features of the primordial density anisotropies over comoving scales of cosmological size can be naturally and simply explained in the framework of standard FRW cosmology, without any need of an earlier stage of inflation in the remote past. The aim of this paper is to complement the results of ref. [1] and show that also the generic statistical features of the primordial density anisotropies over sub-horizon comoving scales can be naturally and simply understood in the standard FRW cosmology.

In short: we claim that in standard FRW cosmology (without inflation) random density anisotropies causally generated at the same instant of recombination have the generic statistical features of the observed primordial anisotropies. Moreover, these characteristic features are indeed derived from the constrain that primordial fluctuations cannot be correlated over

comoving scales longer than the causal horizon at recombination.

Namely, the referred generic statistical features of the observed primordial anisotropies are:

- a) scale invariance over comoving length scales $L \gg H_{eq}^{-1}$ of cosmological size;
- b) a pattern of acoustic (harmonic) peaks over sub-horizon comoving length scales $L \lesssim H_{eq}^{-1}$, located at comoving momenta $k \simeq n \frac{\pi}{H_{eq}}$ with $n = 1, 2, \dots$ a natural number.

In [1] we addressed feature a). We proved there that random density fluctuations causally generated at the instant of recombination, and with comoving wavelength $\lambda \lesssim H_{eq}^{-1}$ shorter than the comoving causal horizon at that instant, produce scale invariant mass anisotropies over comoving volumes of cosmological size $L \gg H_{eq}^{-1}$. Scale invariance means that the variance of random anisotropies in the extensive magnitude M_V (the mass/energy contained in a comoving volume V) is $(\Delta M_V)^2 \sim S$ linearly proportional to the area $S \sim L^2$ of the surface that bounds the region of integration, rather than to its volume $V \sim L^3$. This feature can be intuitively understood in the framework of standard FRW cosmology by noticing that, under standard causal constraints, only random density fluctuations happening at the boundary surface can contribute to fluctuations of the extensive magnitude M_V over cosmologically large comoving volumes. We refer the reader to [1] and [5] for a more detailed explanation of these arguments.

In this paper we address feature b). We start noticing that in standard FRW cosmology it can be consistently (and naturally) claim that the causal horizon H_{eq}^{-1} at the instant of recombination forces the two-point correlation function $f(\vec{x} - \vec{y}) = \langle \rho(\vec{x}) \rho(\vec{y}) \rangle - \langle \rho(\vec{x}) \rangle \langle \rho(\vec{y}) \rangle$, $\vec{x}, \vec{y} \in \mathbf{R}^3$, of the homogeneous and isotropic stochastic density field $\rho(\vec{x})$ to cancel over longer comoving distances,

$$f(\vec{x} - \vec{y}) = 0, \quad \text{if} \quad |\vec{x} - \vec{y}| \geq H_{eq}^{-1}. \quad (1)$$

That is, in standard FRW cosmology causality constrains the two-points correlation function to a compact support, the comoving sphere of radius H_{eq}^{-1} . This condition (1) indeed defines H_{eq}^{-1} : it is the comoving sound causal horizon at recombination. If the two points correlation function has compact support, it accepts a discrete Fourier transform

$$f(|\vec{x} - \vec{y}|) \simeq \left(\frac{\rho_0^2}{2\pi^2} \right) \sum_{n \in \mathbf{N}} \left(\frac{\pi}{H_{eq}^{-1}} \right)^3 \cdot (n^2 \mathcal{P}_n) \frac{\sin \left(\frac{\pi}{H_{eq}^{-1}} n \cdot r \right)}{\left(\frac{\pi}{H_{eq}^{-1}} n \cdot r \right)}, \quad (2)$$

with comoving modes located at $k_n = n \frac{\pi}{H_{eq}^{-1}}$ with $n = 1, 2, \dots$. This discrete power spectrum reproduce the localization of harmonic acoustic peaks in the observable sub-horizon spectrum of primordial cosmological density anisotropies, with a first peak located at $k_1 = \frac{\pi}{H_{eq}^{-1}}$.

In the theoretical framework we have laid the localization of the acoustic peaks is determined by condition (1). These arguments prove that inflation is not necessary to understand the statistical features of the primordial density anisotropies imprinted in the CMB, either their scale invariance over cosmologically large comoving volumes nor the pattern and localization of acoustic peaks over the sub-horizon scales. Furthermore, if our interpretation of the origin of primordial structures is correct it can be used to constrain non-standard cosmologies.

As a last brief comment we wish to notice that the acoustic peaks in the observable power spectrum of temperature anisotropies in the cosmic background radiation are not expected to be sharply located, as expansion (2) could induce to think. They should appear as broad peaks with a width determined by causal effects. In this paper, nevertheless, we will not attempt precise calculations of such effects, because our aim, for the moment, is simply to show that the generic features of the primordial density anisotropies at the instant of recombination clearly correspond to those that should be expected if the anisotropies were causally generated in the framework of standard FRW cosmology, without inflation. In fact, we have shown that the existence in standard FRW cosmology of a comoving causal horizon at the instant of recombination much shorter than all comoving cosmological length scales is not an obstacle for explaining the generic statistical features of the primordial density anisotropies imprinted in the CMB. Just the contrary, the existence of such horizon is the cause of these characteristic features.

[1] D. H. Oaknin, "Spatial correlations of primordial density anisotropies in the standard cosmological model", hep-th/0305068.

- [2] A. Penzias and R.W. Wilson, *Astrophys. J.* **142**, 419 (1965); G. F. Smoot *et al.*, *Astrophys. J.* **396**, L1 (1992); C. L. Bennett *et al.*, *Astrophys. J.* **436**, 423 (1994).
- [3] R.H. Dicke, P.J.E. Peebles, P.G. Roll and D.T. Wilkinson, *Astrophys. J.* **142**, 414 (1965).
- [4] D. N. Spergel *et al.* *Astrophys. J. Suppl.* **148**, 175 (2003); E. Komatsu *et al.*, *Astrophys. J. Suppl.* **148**, 119 (2003); C. B. Netterfield *et al.* [Boomerang Collaboration], *Astrophys. J.* **571**, 604 (2002); C. Pryke, N. W. Halverson, E. M. Leitch, J. Kovac, J. E. Carlstrom, W. L. Holzapfel and M. Dragovan, *Astrophys. J.* **568**, 46 (2002); R. Stompor *et al.*, *Astrophys. J.* **561**, L7 (2001).
- [5] D.H. Oaknin, *Phys. Rev.* **D70**, 103513 (2004).
- [6] See, for example, S. Dodelson, "Coherent phase argument for inflation", hep-ph/0309057.